

## MODEL FODDER PLOT

A model fodder plot with various fodder varieties has been established at this station. The different fodder varieties maintained are as follows:

Fodder varieties	Botanical name
<b><u>Legumes</u></b>	
Hedge lucerne	<i>Desmanthus virgatus</i>
Stylo	<i>Stylosanthes hamata</i>
	<i>Stylosanthes scabra</i>
	<i>Stylosanthes guyanensis</i>
Clitoria/Butterfly Pea	<i>Clitoria ternate</i>
<i>Calopogonium</i>	<i>Calopogonium mucunoides</i>
Lucerne/Alfalfa	<i>Medicago sativa</i>
<i>Desmodium</i>	<i>Desmodium gangeticum</i>
<b><u>Napier hybrid varieties</u></b>	
<i>Cumbu-Napier hybrid (Co4)</i>	<i>Pennisetum purpureum x Pennisetum americanum</i>
<i>Cumbu-Napier hybrid (Co5)</i>	
<i>Australian Red Napier</i>	<i>Pennisetum purpureum cross</i>
<i>APBN1 (Andhra Pradesh Bajra Napier 1)</i>	<i>Pennisetum purpureum cross</i>
<i>Napier grass/Elephant Grass</i>	<i>Pennisetum purpureum</i>
<i>NB21</i>	<i>Pennisetum purpureum x Pennisetum typhoides</i>
<b><u>Forage Pasture grasses</u></b>	
<i>Guinea grass (Bundel variety)</i>	<i>Panicum maximum</i>
<i>Guinea grass (Rivesdale)</i>	<i>Panicum maximum</i>
<i>Guinea grass (GG3)</i>	<i>Panicum maximum</i>
<i>Congo grass</i>	<i>Imperata cylindrica</i>
<i>Buffalo grass</i>	<i>Buchloe dactyloides</i>
<i>Deenanath grass</i>	<i>Pennisetum pedicellatum</i>
<i>Anjan grass</i>	<i>Cenchrus ciliaris</i>
<i>Black Anjan grass</i>	<i>Cenchrus setigerus</i>
<i>Bread Grass</i>	<i>Brachiaria brizantha</i>
<i>Congo signal grass</i>	<i>Brachiaria ruziziensis</i>
<b><u>Multi cutting Sorghum varieties</u></b>	
<i>Fodder sorghum</i>	<i>Co-Fs 29</i>
<i>Fodder sorghum</i>	<i>Co-Fs 31</i>

